

G.C.H. implies existence of many rigid almost free abelian groups*

Rüdiger Göbel
 Fachbereich 6
 Mathematik
 Universität Essen
 45117 Essen, Germany

e-mail:
 R.Goebel@uni-essen.de

Saharon Shelah
 Department of Mathematics
 Hebrew University
 Jerusalem, Israel
 and Rutgers University
 New Brunswick, N. Y., U.S.A.

e-mail:
 Shelah@math.huji.ac.il

February 1, 2008

§1 Introduction

More than thirty years ago A. L. S. Corner [2] proved a very remarkable theorem concerning the existence of “pathological” abelian groups; it is formulated in terms of endomorphism rings.

If A is a countable ring with reduced and torsion-free additive group $A_{\mathbb{Z}}$, then there exists a reduced and countable abelian group G with endomorphism ring $\text{End}_{\mathbb{Z}}G = A$.

This result has been generalized in various directions, e.g. replacing the ground ring $\mathbb{Z}$ by a more general ring and dropping the cardinal restriction (in Corner’s result this is actually $|A| < 2^{\aleph_0}$), see [2], [20], [8], [9], [4], [23].

Other extensions include abelian groups which are not necessarily torsion-free, see Corner, Göbel [4] for references.

Here we are interested in a different strengthening of Corner’s important result: As a byproduct of older constructions of abelian groups G with prescribed endomorphism ring A we have obtained (in a strong sense) almost-free groups provided $A_{\mathbb{Z}}$ is free and provided we are working in Gödel’s universe $V = L$.

*This work is supported by GIF project No. G-0294-081.06/93 of the German-Israeli Foundation for Scientific Research & Development

Let us recall the appropriate definition. An R -module G (of cardinality λ) with $|R| < |G|$ is strongly λ -free if G has a λ -filtration (a continuous chain of submodules $\{G_\nu : \nu < \lambda\}$ with $|G_\nu| < \lambda$ and $\bigcup_{\nu < \lambda} G_\nu = G$) with two additional properties:

- (a) Any G_ν is a free R -module.
- (b) If ν is a successor ordinal and $\nu < \rho < \lambda$, then G_ρ/G_ν is free as well.

We say that $G = \bigcup_{\nu < \lambda} G_\nu$ is a λ -filtration for strong λ -freeness. First results on almost-free groups of cardinality $2^{\aleph_0}$ are in [1].

Because strong λ -freeness is a very natural definition, it can be reformulated in many ways, see Eklof, Mekler [15] for equivalent characterizations and many results on this class of modules. In order to illustrate such results we will restrict ourselves to abelian groups, i.e. $R = \mathbb{Z}$. In $V = L$ we have the following strengthening of Corner's result indicated above.

If A is a ring, $A_{\mathbb{Z}}$ is a free abelian group and $\lambda > |A|$ is any regular, not weakly compact cardinal, then we can find a strongly λ -free abelian group G with $|G| = \lambda$ and $\text{End}_{\mathbb{Z}}G = A$.

This older result [8] grew out of ideas of Eklof, Mekler [14], and Shelah [38]. It is closely related to the work of Jensen [30]. It is only natural to ask if such almost free groups exist without the vehicle of additional set theory. Extending a result of Shelah [36], Eda [11] showed that there is at least some hope.

There exists an $\aleph_1$ -free abelian group G of cardinality $\aleph_1$ with trivial dual $G^* = 0$, see also Corner, Göbel [5]. Recall that $\text{Hom}(G, R) = G^*$ is the dual of an R -module, and G is κ -free for some cardinal κ if any submodule U with $|U| < \kappa$ is contained in a free submodule. In [22] we showed the following stronger result.

If A is a countable ring and $A_{\mathbb{Z}}$ is free, then we can find an $\aleph_1$ -free abelian group G of cardinality $\aleph_1$ with endomorphism ring $\text{End}_{\mathbb{Z}}G = A$.

This result provides a satisfying answer in ZFC, but unfortunately (from the algebraist's point of view) it remains restricted to the world $\aleph_1$.

In fact, assuming ZFC + Martin's axiom, it was shown in [22] that $\aleph_2$ -free groups of size $\aleph_2$ are already always separable (= pure subgroups of products $\mathbb{Z}^{\mathbb{I}}$), hence they must split into various summands. Their endomorphism rings will never be $\mathbb{Z}$ for instance. This brings us to the starting point of this paper.

We want to derive a realization theorem of rings A with $A_{\mathbb{Z}}$ free as $A = \text{End}_{\mathbb{Z}}G$ for strongly $|G|$ -free abelian groups G with $|G| \geq \aleph_1$, using as weak

additional set theoretic conditions as possible. From the result mentioned above we know that Martin's axiom must be excluded. Our results discussed below are surprising in two ways. First of all, we will only need a special case of G.C.H., which ensures the existence of the desired modules. In another paper [43], Shelah and Spasojevic will show by proper forcing arguments that even G.C.H. will not be strong enough for $\text{End } G = \mathbb{Z}$, $|G| = \lambda$ and G λ -free if λ is too large, e.g. if λ is any strongly inaccessible cardinal. Hence we are limited to "small large cardinals". Secondly it turns out that the restriction to G.C.H. forces us to develop novel ideas to get the desired modules. A new Step-Lemma is needed! Here we will also apply combinatorial tools fortunately developed by Shelah [35], [41] a decade ago. The main results of this paper can be summarized as follows.

In order to construct pathological groups we begin with the existence of groups with trivial duals for cardinals $\aleph_n$ ($n \in \omega$). Then we derive results about strongly $\aleph_n$ -free abelian groups of cardinality $\aleph_n$ ($n \in \omega$) with prescribed free, countable endomorphism ring. Finally we use combinatorial results [35], [41] to give similar answers for cardinals $> \aleph_\omega$. As in Magidor and Shelah [31], a paper concerned with the existence of κ -free, non-free abelian groups of cardinality κ , the induction argument breaks down at $\aleph_\omega$. Recall that $\aleph_\omega$ is the first singular cardinal and such groups of cardinality $\aleph_\omega$ do not exist by the well-known Singular Compactness Theorem [34], see also Hill [28] for cardinals cofinal to ω and, for a survey, Eklof and Mekler [15, p. 107].

We will fix a countable commutative ring R with multiplicatively closed subsets S of regular elements containing 1. An R -module G will be called torsion-free (with respect to S) if $sg = 0$ ($s \in S$, $g \in G$) implies $g = 0$. Moreover G is reduced (with respect to S) if $\bigcap_{s \in S} sG = 0$. If A is an R -algebra, and the R -module A_R is R -free (torsion-free or reduced) we will say that the algebra A is free (torsion-free or reduced). We will also fix a countable, free R -algebra A and derive the following results.

Theorem 5.1: *Let (R, S) be as above. If H is a strongly μ -free R -module of regular cardinality μ with trivial dual $H^* = 0$ and $\lambda = 2^\mu = \mu^+$ (the successor cardinal of μ), then there exists a strongly λ -free R -module of cardinality λ with trivial dual.*

As a consequence of (5.1) we obtain the existence of such R -modules of cardinality μ_n ($n \in \omega$) for the n -th successor μ_n of μ provided $2^{\mu_i} = \mu_{i+1}$ for all $i < n$. The existence of H as in (5.1) for $|H| = \aleph_1$ under $2^{\aleph_0} = \aleph_1$ is well-known, see Dugas [7], Shelah [38] or Dugas, Göbel [8], cf. also Eklof, Mekler [15, p. 391]. Hence the existence of the derived strongly $\aleph_n$ -free R -modules of cardinality $\aleph_n$ with trivial dual is immediate (see also (5.2)).

In §6 we will work with a stronger **algebraic** hypothesis, which again is satisfied automatically for $\aleph_1 = 2^{\aleph_0}$ by the papers [8],[38] above. We derive the

Theorem 6.1: *Let (R, S) , $\lambda = 2^\mu = \mu^+$ and μ regular as above. If H is a strongly μ -free R -module of cardinality μ with $\text{End}_R H = A$ for some countable, free R -algebra A , then there exists a strongly λ -free R -module G of cardinality λ with $\text{End}_R G = A$.*

Now the existence of strongly $\aleph_n$ -free R -modules G of cardinality $\aleph_n$ with $\text{End}_R G = A$ follows for each countable, free R -algebra A , provided $\aleph_{i+1} = 2^{\aleph_i}$ for each $i < n$. As one of many consequences we see that under the mild restriction $\aleph_{i+1} = 2^{\aleph_i}$ ($i < n$) there are counterexamples to Kaplansky's test problems for each cardinal $\aleph_n$ ($n \in \omega$) even ones which are strongly $\aleph_n$ -free (see [3] for the required rings). Similarly, we obtain indecomposable, strongly $\aleph_n$ -free groups of cardinality $\aleph_n$. By the result of Shelah, Spasojevic [43] we know that a realization theorem similar to (6.1) and even the existence of modules with trivial dual (5.1) does not follow from G.C.H. for large enough cardinals, e.g. for the first strongly inaccessible cardinal ξ . What happens at smaller cardinals, for instance at $\aleph_{\omega+1}$? The Theorems 6.1 and (5.1) can be extended to $\aleph_{\omega+1}$, hence to $\aleph_{\omega+n}$ ($n \in \omega$) by (5.1) and (6.1), and to certain larger cardinals $< \xi$. We must strengthen the methods from §§3,4 and 5, and this procedure is sketched in §7.

§2 Representation of strongly μ -free modules and the notion of a type of freeness

We fix some notations throughout this paper.

(2.1) *Let R be a commutative ring and $S \subseteq R$ a countable, multiplicatively closed subset of regular elements such that R_R (as R -modules) is reduced and torsion-free (with respect to S), compare §1. We will also assume $1 \in S$ and enumerate $S = \{s_n : n \in \omega\}$ and set $q_n = \prod_{i=1}^n s_i$.*

Our results concern almost free R -modules (mainly for $|R| = \aleph_0$), hence we will refine the notion of almost free and strongly μ -free R -modules from §1:

We say that a strongly μ -free R -module H has freeness type $ftH = u$ if $u = \langle \mu_1, \dots, \mu_n \rangle$ is a descending sequence of regular cardinals μ_i with $\mu_1 = \mu$, $\mu_n = \aleph_0$ and if $H = \bigcup_{\alpha < \mu_1} H_\alpha$ is a μ_1 -filtration for strong μ_1 -freeness with $H_{\alpha+1}/H_\alpha$ either free or of freeness type $\langle \mu_2, \dots, \mu_n \rangle$. The definition is completed by induction if we say that countable groups which are not free are of freeness type $\langle \aleph_0 \rangle$. If H is of freeness type $u = \langle \mu_1, \dots, \mu_n \rangle$, then we can choose a particular filtration of H , which we will call a u -filtration of H . We will write $H = \bigcup_{\alpha_1 < \mu_1} H_{\alpha_1}$, such that $H_{\alpha_1+1}/H_{\alpha_1}$ is either free or of freeness

type $\langle \mu_2, \dots, \mu_n \rangle$ and set $\mathbf{S}_{\mu_1} = \{\alpha \in \mu_1 : H_{\alpha+1}/H_\alpha \text{ is not free} \}$. By inserting $< \mu_1$ members between H_α and $H_{\alpha+1}$ we may assume that $H_{\alpha+1}/H_\alpha \cong R$ if $\alpha \notin \mathbf{S}_{\mu_1}$. Moreover **we assume** $\mathbf{S}_{\mu_1} \subseteq E_{\mu_2} := \{\alpha \in \mu_1, cf\alpha = \mu_2\}$. If $\alpha \in \mathbf{S}_{\mu_1}$, then $H_{\alpha+1}/H_\alpha$ is not free and we proceed similarly. We can find a continuous chain

$$H_\alpha = H_{\alpha_0} \subseteq \dots \subseteq H_{\alpha_\beta} \subseteq \dots \subseteq H_{\alpha_{\mu_2}} = H_{\alpha+1} \quad (\beta < \mu_2)$$

with quotients $H_{\alpha_\beta+1}/H_{\alpha_\beta}$ either free or of type $\langle \mu_3, \dots, \mu_n \rangle$. Recursively we unravel all of $\langle \mu_1, \dots, \mu_n \rangle$ and let $u^i = \mu_1 \times \dots \times \mu_i$ ($i = 1, \dots, n$). We obtain a refined, continuous chain $\{H_\alpha, \alpha \in u^i\}$ of free R -modules H_α with $H_\beta/H_{\alpha+1}$ free for $\beta \geq \alpha + 1$. Moreover $H_{\alpha+1}/H_\alpha$ is either $\cong R$ or of type $\langle \mu_{i+1}, \dots, \mu_n \rangle$. The set u^i is ordered lexicographically and each $\alpha \in u^i$ is of the form $\alpha = \alpha_1^\wedge \dots^\wedge \alpha_i$ with $\alpha_j < \mu_j$. As usual, $\alpha + 1$ will denote the successor of α in u^i which is $\alpha + 1 = \alpha_1^\wedge \dots^\wedge \alpha_i + 1$, and 0 denotes the initial element of u^i .

Moreover, again refining the chain $\{H_\alpha : \alpha \in u^n\}$, we may assume that $H_{\alpha+1}/H_\alpha \subseteq S^{-1}R$ for each $\alpha \in u^n$ and $H_0 = 0$. We will identify u^n with μ in §§3 – 6 and write for $\mathbf{S} = \{\alpha \in \mu : H_{\alpha+1}/H_\alpha \text{ is not free} \}$, hence

$$(2.2) \begin{cases} \text{either } H_{\alpha+1} = H_\alpha \oplus t_{\alpha 0}R, \text{ if } \alpha \notin \mathbf{S} \text{ or } H_{\alpha+1} = \langle H_\alpha, t_{\alpha k} : k \in \omega \rangle \\ \text{is a free } R\text{-module with } q_{\alpha k} t_{\alpha k+1} = t_{\alpha k} + \sum_{j < k_\alpha} t_{\alpha_j k_j} \text{ where} \\ q_{\alpha k} \in S, t_{\alpha_j k_j} \in H_{\alpha_j+1} \text{ for } \alpha_j < \alpha \text{ and } k < \omega. \end{cases}$$

Remark: The reader not interested in particular freeness-type or in modules of rank $\aleph_{\omega+1}$ may choose (2.2) immediately from $\{H_\alpha, \alpha \in u^1\}$. Recall that H is uniquely determined (up to isomorphism) by these generators and relations (as follows by a transfinite induction on $\alpha < \mu$). Inductively we will also use particular free resolutions.

Definition 2.3: If $u = \langle \mu_1, \dots, \mu_n \rangle$ is a (finite) descending sequence of regular cardinals, then we say that the R -module M has a u -resolution for $\kappa \in u$, if $\mu_1 < \mu$ and the following holds:

- (1) M is strongly μ -free of cardinality μ .
- (2) We have a free resolution $M \cong F/K$.
- (3) $K = \bigoplus_{k \in \kappa} K_k$ and F, K_k are free R -modules of rank μ .
- (4) K_k has a basis $\{x_{\alpha k} : \alpha < \mu\}$ and if $K_{\beta k} = \langle x_{\alpha j} : j < k \text{ or } \alpha < \beta \rangle$ for $(k < \kappa) (\beta < \mu)$, then $F/K_{\beta k}$ is free.

Sometimes R in (2.3) will be replaced by an R -algebra A , provided M is a strongly μ -free A -module.

§3 A Step-Lemma and a Freeness-Proposition

The μ - $\aleph_0$ -Step-Lemma 3.1: *Let H be a strongly μ -free R -module of cardinality μ and $H^* = 0$. Moreover, let $F = \bigoplus_{n \in \omega} K_n$ be a direct sum of free modules K_n of rank μ . Choose a partial basis $x_{\alpha n}$ ($\alpha < \mu$) for each K_n , let $K'_n = \langle x_{\alpha n} : \alpha < \mu \rangle$ and let $f \in K_0$ be any pure element with K_0/fR free. Then we can find two extensions $F^i \supset F$ ($i = 0, 1$) with the following properties.*

- (1) *If $K_{\beta n} = \langle x_{\alpha i} : i < n \text{ or } \alpha < \beta \rangle = \bigoplus_{i < n} K'_i \oplus \langle x_{\alpha i} : i \geq n, \alpha < \beta \rangle$ then $F^i/K_{\beta n}$ is free R -module for each $n \in \omega$, $\beta < \mu$.*
- (2) *$F^i/F \cong H$ ($i = 0, 1$).*
- (3) *If $\varphi \in F^*$ extends to both $\varphi^i \in F^{i*}$ ($i = 0, 1$), then $f\varphi = 0$.*

Remarks: It follows from (1) that F^i is free, because $K_{\beta n}$ is free as well. The particular case $\beta = 0$ with $K_{0n} = \bigoplus_{i < n} K_i$ looks more familiar in connection with older Step-Lemmas, but we need more – that is stated in (1). Conditions (1) and (2) say that H has an $\langle \aleph_0 \rangle$ -resolution for $\aleph_0$ as defined in (2.3).

Later on we will need a more general Step-Lemma as well, in order to deal with abelian groups of cardinality $\geq \aleph_{\omega+1}$. This however is postponed to Section 7.

Proof. Recall our representation (2.2) of H . We have

$$(i) \begin{cases} H_{\alpha+1} &= \langle H_\alpha, t_{\alpha n} : n \in \omega \rangle \text{ with} \\ q_{\alpha n} t_{\alpha n+1} &= t_{\alpha n} + \sum_{j < n_\alpha} t_{\alpha_j n_j} \text{ if } \alpha \in \mathbf{S} \text{ and} \\ H_{\alpha+1} &= H_\alpha \oplus t_\alpha R \text{ if } \alpha \in \mu \setminus \mathbf{S} \end{cases}$$

where $q_{\alpha n} \in S$, $t_{\alpha_j n_j} \in H_{\alpha_j+1}$ for $\alpha_j < \alpha$. Let $C^i = \bigoplus_{\alpha \in \mathbf{S}} \bigoplus_{n \in \omega} s_{\alpha n}^i R \oplus \bigoplus_{\alpha \in \mu \setminus \mathbf{S}} s_\alpha^i R$ be freely generated by elements $s_{\alpha n}^i$ and s_α^i for $i = 0, 1$. Next we choose two submodules $N^i \subseteq F \oplus C^i$ by generators

$$N^0 = \langle q_{\alpha n} s_{\alpha n+1}^0 - s_{\alpha n}^0 - \sum_{j < n_\alpha} s_{\alpha_j n_j}^0 - x_{\alpha n} : \alpha \in \mathbf{S}, n \in \omega \rangle$$

and

$$N^1 = \langle q_{\alpha n} s_{\alpha n+1}^1 - s_{\alpha n}^1 - \sum_{j < n_\alpha} s_{\alpha_j n_j}^1 - x_{\alpha n} - f : \alpha \in \mathbf{S}, n \in \omega \rangle,$$

where $f \in K_0$ is taken from the hypothesis of the Lemma. If $F^i = F \oplus C^i/N^i$, then we identify $x \in F \oplus C^i$ with $x + N^i$ in F^i , hence $F \subset F^i$ ($i = 0, 1$) and the following relations hold in F^i by choice of N^i :

$$(r^i) \quad q_{\alpha n} s_{\alpha n+1}^i = s_{\alpha n}^i + \sum_{j < n_\alpha} s_{\alpha_j n_j}^i + x_{\alpha n} + f \cdot \delta_{1i} \quad (i = 0, 1)$$

where $\delta_{11} = 1$ and $\delta_{10} = 0$.

In order to show (2), we consider the maps $\psi_i : F^i \rightarrow H$ defined by $\psi_i|_F = 0$ and $s_{\alpha n}^i \psi_i = t_{\alpha n}$ for all $\alpha \in \mathbf{S}$, $n \in \omega$, $s_{\alpha}^i \psi_i = t_{\alpha}$ for $\alpha \in \mu \setminus \mathbf{S}$ and $i = 0, 1$. These maps preserve the relations (r^i) and (i) , hence ψ_i can be extended to an R -homomorphism $\psi_i : F^i \rightarrow H$. Its kernel is clearly F , ψ is epic and (2) follows.

We have a decomposition into free summands $K'_n \oplus L_n = K_n$ where $K'_n = \langle x_{\alpha n} : \alpha < \mu \rangle$, and L_n comes from the complement of the partial basis, hence

$$(ii) \begin{cases} F &= \oplus_{m < n} K_m \oplus \oplus_{m \geq n} L_m \oplus \langle x_{\alpha m} : \alpha < \mu, m \geq n \rangle \text{ and} \\ F^0 &= \oplus_{m < n} K_m \oplus \oplus_{m \geq n} L_m \oplus C \text{ where} \\ C &= \langle x_{\alpha m}, s_{\alpha k}^0, s_{\beta}^0 : \alpha \in \mathbf{S}, \beta \in \mu \setminus \mathbf{S}, m \geq n, k \in \omega \rangle \end{cases}$$

Using (r^0) we see that the $x_{\alpha m}$'s can be expressed by $s_{\alpha k}^0$'s, hence C is also generated by

$$C = \langle s_{\alpha k}^0, s_{\beta}^0 : \alpha \in \mathbf{S}, \beta \in \mu \setminus \mathbf{S}, k < \omega \rangle$$

If $i = 0$ is replaced by $i = 1$ we note that $f \in K_0$ and a similar submodule C is a new complement satisfying (ii). In order to show (1), we consider the case $i = 0$ only. Let $- : F^0 \rightarrow F^0/K_{\beta n}$ denote the canonical epimorphism. The first step in proving that $\overline{F}^0$ is free is

$$(iii) \begin{cases} \overline{C}_{n\beta} := \langle \overline{s}_{\alpha k}^0, \overline{s}_{\alpha'}^0 : \alpha' \in \mu \setminus \mathbf{S} \text{ and } \alpha \in \mathbf{S} \text{ such that} \\ k < n \text{ or } k \geq n \text{ and } \alpha', \alpha < \beta \rangle \subseteq \overline{C} \text{ is freely generated.} \end{cases}$$

Note that

$C_{\gamma} := \langle \overline{s}_{\alpha k}^0, \overline{s}_{\alpha'}^0 : \alpha \in \mathbf{S}, \alpha' \in \mu \setminus \mathbf{S} \text{ with } k < n, \alpha', \alpha < \gamma \text{ or } k \geq n, \alpha', \alpha < \beta \rangle$
 $(\beta \leq \gamma < \mu)$ is an ascending, continuous chain of submodules of $\overline{C}_{n\beta}$ with $\overline{C}_{n\beta} = \bigcup_{\gamma < \mu} C_{\gamma}$. Because $|C_{\gamma}| \leq |\gamma| + |\beta| + \aleph_0 < \mu$ and $C_{\gamma} \subseteq \overline{F}^0$ is strongly μ -free by (2), as an extension of a free module by a strongly μ -free module, each C_{γ} is free as well. Now (iii) will follow if we show

$$C_{\gamma+1}/C_{\gamma} \cong \langle \overline{s}_{\gamma n-1}^0 + C_{\gamma} \rangle \quad (\cong R)$$

for each $\gamma \geq \beta$. Observe that this is trivial if $\gamma \notin \mathbf{S}$, hence we assume $\gamma \in \mathbf{S}$. For simplicity we will ignore the elements s_{α}^i ($\alpha \notin \mathbf{S}$)! The relation (r^0) under $-$ become

$$q_{\gamma k} \overline{s}_{\gamma k+1}^0 = \overline{s}_{\gamma k}^0 + \sum_{j < n_{\gamma}} \overline{s}_{\gamma_j n_j}^0,$$

as in (i), and $C_{\gamma+1}/C_{\gamma} = \langle \overline{s}_{\gamma k}^0 + C_{\gamma} : k < n \rangle$. For large enough $q \in S$ we get $q \overline{s}_{\gamma n-1}^0 = q' \overline{s}_{\gamma 0}^0 + \sum_j q_j \overline{s}_{\beta_j 0}$ and $\sum_j q_j \overline{s}_{\beta_j 0}$ is in C_{γ} , $q'|q$. Hence the n generators above can be replaced by $C_{\gamma+1}/C_{\gamma} = \langle \overline{s}_{\gamma n-1}^0 + C_{\gamma} \rangle$ and (iii) holds.

Next we show that

$$(iv) \begin{cases} \overline{C}_{n\beta} \oplus \overline{C}^{n\beta} = \overline{C} \text{ where } \overline{C}^{n\beta} = \langle \overline{s}_{\alpha k}^0 : \beta \leq \alpha < \mu, k \geq n \rangle \\ \text{is freely generated by the } \overline{s}_{\alpha k}^0 s. \end{cases}$$

Clearly $\overline{C}$ is generated by $\overline{C}_{n\beta}$ and $\overline{C}^{n\beta}$.

Any linear combination $s = \sum r_{\alpha k} \overline{s}_{\alpha k}^0 \in \overline{C}^{n\beta}$ may be multiplied by a large enough $q \in S$ such that the summand $qr_{\alpha k} \overline{s}_{\alpha k}^0$ by (r^0) becomes a sum with summand $q'r_{\alpha k} x_{\alpha k}$. Now $s \in \overline{C}_{n\beta}$ is only possible if all the independent terms $q'r_{\alpha k} x_{\alpha k}$ ($\alpha \geq \beta$, $k \geq n$) vanish, hence $r_{\alpha k} = 0$ and $s = 0$. The sum (iv) must be direct. The freeness of $\overline{C}^{n\beta}$ follows similarly, again using independence of the $x_{\alpha k}$ under $-$, and (iv) follows.

The quotient $\overline{F}^0 / \overline{C}_{n\beta} \cong \bigoplus_{m \geq n} L_m \oplus \overline{C}^{n\beta}$ is free by (iv) and $\overline{F}^0$ is free by (iii) and (1) is shown for $i = 0$.

In order to show (3) we consider any $\varphi \in F^*$ extending to $\varphi^i \in F^{i*}$ for $i = 0, 1$ and recall that $f \in F_0 \subseteq F$ is a prescribed basic element by hypothesis.

First we consider the pushout D of $F \subset F^i$, which is

$$D = F^0 \oplus F^1 / \triangle \text{ with } \triangle = \{(g, -g) : g \in F\} \cong F.$$

We also have a pushout extension $\varphi^i \subset \Phi \in D^*$ ($i = 0, 1$) by the pushout properties and $\varphi \subseteq \varphi^i$.

If $\overline{g} = (g, 0) + \triangle$, $\overline{g} = (0, g) + \triangle$ for $g \in F^0$ and $g \in F^1$ respectively, then we have canonical identifications $\overline{F}^0 + \overline{F}^1 = D$ with

$$\overline{F} = \overline{F}^0 \cap \overline{F}^1 = \{(g, 0) + \triangle : g \in F\} = \{(0, g) + \triangle : g \in F\}.$$

We omit “ $-$ ” in the following. Let $U =$

$\langle F, d_{\alpha n}, d_{\alpha'} : \alpha \in S, n < \omega, \alpha' \in \mu \setminus S \rangle \subseteq D$ for $d_{\alpha n} = s_{\alpha n}^0 - s_{\alpha n}^1$, $d_{\alpha} = s_{\alpha}^0 - s_{\alpha}^1$. Observe that

$$U + F^i = F^0 + F^1 = D \text{ and } U \cap F^i = F^0 \cap F^1 = F.$$

The desired relations from (r^i) are

$$(v) \left\{ \begin{array}{l} q_{\alpha n} d_{\alpha n+1} = d_{\alpha n} + \sum_{i < n_{\alpha}} d_{\alpha_i n_i} + f. \\ \text{The map } t_{\alpha n} \rightarrow d_{\alpha n} + \langle f \rangle \text{ induces } V / \langle f \rangle \cong H, \text{ with} \\ V = \langle f, d_{\alpha n}, d_{\alpha'} : \alpha \in S, n \in \omega, \alpha' \in \mu \setminus S \rangle. \end{array} \right.$$

In particular, $U/F \cong H$. Let $f\varphi = r \in R$ and consider the new homomorphism $\psi = \Phi - r : D \rightarrow R$.

Now we have $f\psi = f(\Phi - r) = 0$ and the relations (v) become

$$q_{\alpha n} (d_{\alpha n+1} \psi) = (d_{\alpha n} \psi) + \sum_{i < n_{\alpha}} (d_{\alpha_i n_i} \psi)$$

and $t_{\alpha n} \xrightarrow{\eta} d_{\alpha n} \psi$, $f\psi = 0$, induces an epimorphism

$$\eta : H \rightarrow V\psi$$

and $H^* = 0$ forces $V\psi = 0$. Hence $\Phi|V$ is multiplication by r , which by cardinality reasons is only possible for $r = 0$. We conclude $f\varphi = 0$.

The following Freeness-Proposition is a special case of a more general Freeness-Proposition. In order to make the idea of the construction more transparent, we postpone again the extension of (3.2) to Section 7. The more general Freeness-Proposition 7.2 involves a few more technical difficulties, due to more complicated notation. The Proposition will ensure that continuous chains of free submodules in our construction remain free at limit ordinals below μ^+ .

The ω -Freeness-Proposition 3.2: *Let μ be a regular cardinal and $\delta < \mu^+$ be an ordinal with $\{G_i : i \leq \delta\}$ an ascending, continuous chain of submodules with the following properties for all $i < \delta$.*

- (a) G_i is free of rank μ .
- (b) G_{i+1}/G_i is strongly μ -free and let $\mathbf{S} = \{i < \delta : G_{i+1}/G_i \text{ is not free}\}$
- (c) There exist $G_i \subseteq H_i \subseteq G_{i+1}$ and H_i^0 a free summand of G_i with free quotient such that $H_i = G_i \oplus_{H_i^0} H_i^1$ is a pushout with $H_i^0 \subseteq H_i^1$ free modules of rank μ .
- (d) $G_{i+1} = H_i \oplus K_i$ with K_i freely generated by $x_{\alpha i}$ ($\alpha < \mu$).
- (e) If $i \notin \mathbf{S}$, then $G_i = H_i$ and $H_i^1 = H_i^0$ (hence $G_{i+1} = G_i \oplus K_i$).
- (f) If $i \in \mathbf{S}$, then $\text{cf}(i) = \omega$. Moreover
 - (i) there exists $a_i \subseteq i$ of order type ω with $a_i \cap \mathbf{S} = \emptyset$ and $\sup a_i = i$.
 - (ii) there exist $A_{ij} \subseteq \mu$ for all $j \in a_i$ such that $A_{ij} \cap A_{ij'} = \emptyset$ for $j \neq j'$ and $A_{ij} \cap A_{i'j'}$ is bounded in μ for $(ij) \neq (i'j')$.
 - (iii) Let $K_j^\omega = \langle x_{\alpha j} : \alpha \in A_{ij} \rangle$, $H_i^0 = \bigoplus_{j \in a_i} K_j^\omega$. If $k \in a_i$, $\beta < \mu$ and $K_{\beta k} = \bigoplus_{j \in a_i}^{j < k} K_j^\omega \oplus \langle x_{\alpha j} : j \in a_i, j \geq k, \alpha < \beta \rangle$, then $H_i^1/K_{\beta k}$ is free.

It follows that G_δ is free as well.

Proof. Enumerate $\mathbf{S} = \{i_\epsilon : \epsilon < \sigma\}$ for some $\sigma \leq \mu$. By induction on $\epsilon < \sigma$ choose ordinals $\alpha_\epsilon < \mu$ and suppose $\alpha_0, \dots, \alpha_\nu$ are defined for all $\nu < \epsilon$. Define α_ϵ subject to the following condition

$$(+) \text{ For all } \nu < \epsilon, j \in a_{i_\nu} \cap a_{i_\epsilon} \text{ we have } A_{i_\epsilon j} \cap A_{i_\nu j} = \emptyset.$$

This can be arranged: we collect all troublemakers at stage ϵ , which is a set $M = \bigcup_{\nu < \epsilon} \bigcup_{j \in a_{i_\nu} \cap a_{i_\epsilon}} (A_{i_\nu j} \cap A_{i_\epsilon j}) \subseteq \mu$, and observe that $A_{i_\nu j} \cap A_{i_\epsilon j}$ is bounded below μ by (f)(ii), μ is regular and $|\epsilon| < \mu$. Hence M is bounded by some $\alpha_\epsilon < \mu$ and (+) follows. Rename $\alpha_\epsilon = \alpha_{i_\epsilon}$. Then (+) can be restated as

$$(+) \text{ For all } i \neq t \in \mathbf{S}, j \in a_i \cap a_t \text{ we have } A_{tj} \setminus \alpha_t \cap A_{ij} \setminus \alpha_i = \emptyset.$$

If $i \in \mathbf{S}$, choose any fixed $\beta_i < \mu$ such that $\alpha_i < \beta_i$ and observe that $H_i^0 = K_i^* \oplus C_i$ by (f)(iii) for $k = \min a_i$, $\beta = \beta_i$ and $K_{k\beta} = K_i^* := \langle x_{\alpha j} : j \in a_i, \alpha < \beta_i \rangle$. Now we are able to collect a basis of G_δ : Choose all basis elements $x_{\alpha j}$ not involved in extension of G_i at $i \in \mathbf{S}$, and we have the set

$$B = \{x_{\alpha j} : j < \delta, \text{ there is no } i \in \mathbf{S} \text{ with } j \in a_i, \alpha \in A_{ij} \text{ and } \alpha > \beta_i\}.$$

Moreover $C = \bigoplus_{i \in \mathbf{S}} C_i$ is direct because of (+), and each C_i is free by (f)(iii). We may collect its basis B_C . Finally choose all basis elements $x_{\alpha j}$ ($\alpha < \beta_i$) which are used for the distinct K_i^* , say B' . Then $B \cup B_C \cup B'$ is a basis for G_δ .

§4 The prediction principles

We will use prediction principles that hold under G.C.H. First recall some well-known definitions, tailored for our applications. Let κ be a regular cardinal and $\theta = \{A_\nu : \nu < \kappa\}$ be a κ -filtration of $A = \bigcup_{\nu < \kappa} A_\nu$, i.e. θ is an ascending continuous chain with $A_0 = 0$ and $|A_\nu| < \kappa$ for all $\nu < \kappa$. Let $E \subseteq \kappa$. Then $\diamond_\kappa(E)$ holds if there is a family $\{S_\nu \subseteq A_\nu : \nu \in E\}$ such that for all $X \subseteq A$ the set $\{\nu \in E : X \cap A_\nu = S_\nu\}$ is stationary in κ . Besides this diamond principle, we also consider the weak diamond $\Phi_\kappa(E)$. If $P_\nu : \mathcal{P}(A_\nu) \rightarrow 2 = \{0, 1\}$ ($\nu \in E$) is a given partition of subsets of A_ν , then $\Phi_\kappa(E)$ provides a prediction function $\varphi : E \rightarrow 2$ such that for all $X \subseteq A$ the set $\{\nu \in E : P_\nu(X \cap A_\nu) = \varphi(\nu)\}$ is stationary in κ . In this case E is called non-small. While $\diamond_\kappa(E)$ holds in $V = L$ for all stationary subsets E of regular, not weakly compact cardinals by Jensen [30], prediction principles are quite often valid under weaker assumptions, cf. Eklof, Mekler [15, pp.175-178], Shelah [35, p.376, Theorem 32]. See also Gregory [24].

Proposition 4.1 (Shelah [35]): *Suppose $\lambda = 2^\nu = \mu^+$ and for some regular cardinal $\kappa < \mu$ either*

- (i) $\mu^\kappa = \mu$ or
- (ii) μ is singular with $cf\mu \neq \kappa$ and $|\delta|^\kappa < \mu$ for all $\delta < \mu$

Then $\diamond_\lambda(E_\kappa)$ holds, where $E_\kappa = \{\alpha < \lambda : cf\alpha = \kappa\}$.

In particular we have a

Corollary 4.2: *Assume G.C.H. and μ is some regular cardinal. Then $\diamond_\mu E_\kappa$ holds for any regular cardinal $\kappa < \mu$.*

In the first application we will use $\Phi_\mu(E_{\aleph_0})$, a weak consequence of (4.2). Later we will replace $\kappa = \aleph_0$ by some regular $\kappa > \aleph_0$ (cf. §7). Moreover we will use

Proposition 4.3: *If $\lambda = \mu^+$ and $E \subseteq \lambda$ is non-small, then we can decompose $E = \coprod_{\beta < \kappa} E_\beta$ and each E_β is non-small as well.*

Proof. See Eklof and Mekler [15, p.144].

§5 Construction of modules with trivial dual

We apply the μ - $\aleph_0$ -Step-Lemma 3.1, the ω -Freeness-Proposition 3.2 and combinatorial ideas from §4. The arguments are however partly standard (for working in L), so we keep them short. The reader may consult [15], [38] or [8] for details.

Let $\lambda = \mu^+ = 2^\mu$ for some regular cardinal μ , and let $E = \{\alpha \in \lambda, cf\alpha = \omega\}$. Recall that E is non-small by (4.1), for instance. Hence we assume $\Phi_\lambda(E)$, the weak diamond and decompose $E = \coprod_{\beta < \lambda} E_\beta$ into non-small sets E_β , cf. (4.3). For each $i \in E$ we also fix $a_i = \{i_n : n \in \omega\}$, a strictly increasing sequence $i_n < i$ of successor ordinals i_n with $\sup a_i = i$, hence $E \cap a_i = \emptyset$. For each $j \in a_i$ we find $A_{ij} \subseteq \mu$ with:

- (a) A_{ij} unbounded in μ and $A_{ij} \cap A_{i'j'}$ bounded in μ if $(ij) \neq (i'j')$ and
- (b) $A_{ij} \cap A_{ij'} = \emptyset$ for $j \neq j' \in a_i$.

Choose a set of $2^\mu = \lambda$ subsets $X', X \subseteq \mu$ with $|X \cap X'| < \mu$ for distinct pairs X, X' . Decompose each X into ω subsets of cardinal μ to define the A_{ij} 's. Hence (b) follows and (a) holds because μ is regular. Suppose now that H is a given strongly μ -free R -module with $|H| = \mu$ and trivial dual $H^* = \text{Hom}(H, R) = 0$.

We want to derive the existence of a strongly λ -free R -module G of cardinality λ such that $G^* = 0$.

We have to fix the “prediction” and choose a λ -filtration $\{G_i : i < \lambda\}$ such that $G_0 = 0$ and $|G_i| = |G_{i+1} \setminus G_i| = \mu$ for all $i < \lambda$. Let $\{g_\nu : \nu < \lambda\} = G$ be an enumeration of $G = \bigcup_{i < \lambda} G_i$ such that $g_\beta \in G_i$ for all $i \in E_\beta$. We define partitions $P_i^\beta : \mathcal{P}(G_i \times R) \rightarrow 2$ for all $\beta < i$, $i \in E_\beta$ predicting homomorphisms in G_i^* .

Formally we also must predict an R -module structure on G_i , hence $G_i \times R$ must be replaced by $G_i \times G_i \times \cdots \times R$ to take care of this; however we will ignore this for simplicity; compare [8] or [19, p.284].

If $X \subseteq G_i \times R$, then define $P_i^\beta(X) = 0$ if the following holds.

$$(1) \left\{ \begin{array}{l} G_i, G_j, G_i/G_j \text{ are free } R\text{-modules for } j \in a_i \text{ and } g_\beta \in G_{j_0}, \text{ with } G_{j_0}/g_\beta R \\ \text{free, say } j_0 = \min a_i. \text{ Moreover } G_{j+1} = G_j \oplus K_j \text{ with } K_j \\ \text{freely generated by some } x_{\alpha j} \ (\alpha < \mu). \end{array} \right.$$

$$(2) \left\{ \begin{array}{l} X \text{ is (as a graph) a homomorphism } h_X \in G_i^*. \text{ If } K_j^\omega = \langle x_{\alpha j} : \alpha \in A_{ij} \rangle \\ (j \in a_i) \text{ and if we identify } (G_i, \oplus_{j < n} K_j^\omega, g_\beta, h_X : n \in \omega) \text{ with} \\ (F, \oplus_{j < n} K'_j, f, \varphi : n \in \omega) \text{ in (3.1), then we} \\ \textbf{require} \text{ that } \varphi \text{ does not extend to } F^{0*}. \end{array} \right.$$

From $\Phi_\lambda(E_\beta)$ we have prediction functions $\varphi_\beta : E_\beta \rightarrow 2$ such that

$$\mathcal{X}_\beta(X) := \{i \in E_\beta, P_i^\beta(X \cap G_i \times R) = \varphi_\beta(i)\}$$

is stationary in λ for all $X \subseteq G$, $\beta < \lambda$. The structure of the R -module $G = \bigcup_{i < \lambda} G_i$ is given inductively on the continuous chain $\{G_i : i < \lambda\}$ subject to the following conditions.

(i) G_i is a free R -module for any $i \in \lambda$.

$$(ii) \left\{ \begin{array}{l} G_{i+1} = H_i \oplus K_i \text{ where } K_i \text{ is freely generated by elements} \\ x_{\alpha i} \ (\alpha < \mu), \ G_i \subseteq H_i \subseteq G_{i+1}, \ G_{i+1}/G_i \text{ is strongly } \mu\text{-free} \\ \text{and } H_i = G_i \oplus_{H_i^0} H_i^1 \text{ is a pushout for some free summand} \\ H_i^0 = G_i \cap H_i^1 \text{ of } G_i \text{ with free quotient and } H_i^0 \subseteq H_i^1 \text{ all} \\ \text{free of rank } \mu. \end{array} \right.$$

$$(iii) \left\{ \begin{array}{l} \text{If } i \in E_\beta \text{ and (1) holds, then we choose } H_i \text{ as an extension} \\ F^{\varphi_\beta(i)} \text{ given by the identification } (F, \oplus_{j < n} K'_j, f, \varphi : n \in \omega) \\ \text{in (3.1) with } (G_i, \oplus_{j < n} K_j^\omega, g_\beta, h_X : n \in \omega). \end{array} \right.$$

Let $H_i = G_i$ otherwise. Hence G is defined.

Note that $G_\delta = \bigcup_{i < \delta} G_i$ is a free R -module for limit ordinals $\delta < \lambda$ as follows from (3.2). We conclude that G is a strongly λ -free R -module of cardinality λ by (3.1) and the construction above.

If $\psi \in G^*$ and $\psi \neq 0$, then $C = \{i < \lambda : \psi|G_i \neq 0\}$ is a cub. Moreover $f\psi \neq 0$ for some $f \in G$ with G_i/fR free for large enough $i \in C$, and $f = g_\beta \in G_i$ for some $\beta < \lambda$. Choose an ordinal $i \in C \cap \mathcal{X}_\beta(\psi)$ and observe that $\psi|G_i$ extends to G_{i+1} , hence $P_i^\beta(\psi|G_i) = 1$ and $G_{i+1} = F^1$ as in the Step-Lemma 3.1. However $P_i^\beta(\psi|G_i) = 1$ and (2) tell us that $\psi|G_i$ extends to F^0 as in (3.1) as well. Therefore $f\psi = 0$ by (3.1), a contradiction. Hence $G^* = 0$ follows.

We conclude the following results.

Theorem 5.1: *Let R be a countable commutative ring as in §2. If H is a strongly μ -free R -module of regular cardinality μ with trivial dual $H^* = 0$ and $\lambda = 2^\mu = \mu^+$, then there exists a strongly λ -free R -module of cardinality λ with trivial dual.*

Corollary 5.2: *If $2^{\aleph_i} = \aleph_{i+1}$ for $i = 0, \dots, n-1$, then there exists a strongly $\aleph_n$ -free abelian group of cardinality $\aleph_n$ with trivial dual.*

The corollary follows by induction. If $i = 0$, then any subgroup of $\mathbb{Q}$ different from $\mathbb{Z}$ serves as H in (5.1). From (5.1) we obtain a strongly $\aleph_1$ -free abelian group of cardinality $\aleph_1$ with trivial dual, and so on.

§6 Endo-rigid modules

The following result establishes the existence of R -modules with endomorphism ring R , the so-called rigid R -modules. Such modules are known in ZFC, see e.g. Corner [2] or [20], [21], [40]. However we will also assume that the R -modules in question are almost free. A result in $ZFC + MA$ (= Martin's axiom) shows that extra axioms are needed, because such modules turn out to be separable under $ZFC + MA$, see Göbel, Shelah [22]. Assuming $V = L$, almost free modules were constructed in Dugas [7], Shelah [38] and Dugas, Göbel [8] several years ago. Here we have to work harder to derive the existence under ZFC and a weak form of G.C.H.

Theorem 6.1: *Let R be a countable, commutative ring as in (2.1) and assume $\lambda = 2^\mu = \mu^+$ for some regular cardinal μ . Suppose there exists a strongly μ -free R -module H of cardinality μ with $\text{End } H = R$. Then we can find a strongly λ -free R -module G of cardinality λ with $\text{End } G = R$.*

The proof is – as in (5.1) – divided into three parts, a Step-Lemma similar to (3.1), however with crucial, not-so-obvious changes. The construction of G and the actual proof of the claims in (6.1) are similar to §5. We will keep the proof short. However, the reader familiar with §§3-5 will follow the arguments easily. We begin with the Step-Lemma.

The μ - $\aleph_0$ -Step-Lemma 6.2: *Let R be as in §2 and let H be a strongly μ -free R -module of cardinality μ with $\text{End } H = R$. Moreover, let $F = \bigoplus_{n \in \omega} K_n$ be a direct sum of free modules K_n of rank μ . Choose a partial basis $x_{\alpha n}$ ($\alpha < \mu$) for each K_n , let $f \in K_0$ and $h \in H$ such that K_0/fR is free and H/hR is strongly*

μ -free. Then we can find two extensions $F \subseteq F^i$ ($i = 0, 1$) with the following properties.

- (1) If $K_{\beta n} = \langle x_{\alpha i} : i < n \text{ or } \alpha < \beta \rangle$, then $F^i/K_{\beta n}$ is a free R -module for each $n \in \omega$, $\beta < \mu$.
- (2) $F^i/F \cong H/hR$ is strongly μ -free.
- (3) If $\varphi \in \text{End } F$ extends to both $\varphi^i \in \text{End } F^i$ ($i = 0, 1$), then $f\varphi \in fR$.

Proof. Let $H' = H/hR$ and choose a μ -filtration $H = \bigcup_{i < \mu} H_i$ for strong μ -freeness with $h \in H_0$. Since H/hR is strongly μ -free, also $H'_i = H_i/hR$ is free and $\{H'_i : i < \mu\}$ constitutes a μ -filtration of H' for strong μ -freeness.

Let $\mathbf{S} = \{\alpha < \mu \text{ with } H_{\alpha+1}/H_\alpha \text{ not free}\}$. Next we apply the representation (2.2) to H' and obtain

$$(*) \left\{ \begin{array}{l} \text{a continuous chain of free submodules } H_\alpha \text{ } (\alpha < \mu) \text{ of} \\ H = \bigcup_{\alpha < \mu} H_\alpha \text{ such that } H_0 = hR \text{ and } H_{\alpha+1} = H_\alpha \oplus t_\alpha R \\ \text{if } \alpha \notin \mathbf{S} \text{ or } H_{\alpha+1} = \langle H_\alpha, t_{\alpha n} : n < \omega \rangle \text{ with} \\ q_{\alpha n} t_{\alpha n+1} = t_{\alpha n} + \sum_{j < n_\alpha} t_{\alpha_j n_j} + r_{\alpha n} h \text{ for} \\ q_{\alpha n}, r_{\alpha n} \in S, t_{\alpha_j n_j} \in H_{\alpha_j+1} \text{ with } \alpha_j < \alpha, n < \omega. \end{array} \right.$$

Similar to (3.1), but identifying f and h , we define two extensions F^i of F for $i = 0, 1$. Let $C^i = \bigoplus_{\alpha \in \mathbf{S}} \bigoplus_{n < \omega} s_{\alpha n}^i R \oplus \bigoplus_{\alpha \in \mu \setminus \mathbf{S}} s_{\alpha}^i R$ be freely generated and $N^i \subseteq F \oplus C^i$ such that

$$\begin{aligned} N^0 &= \langle q_{\alpha n} s_{\alpha n+1}^0 - s_{\alpha n}^0 - \sum_{j < n_\alpha} s_{\alpha_j n_j}^0 - x_{\alpha n} : \alpha \in \mathbf{S}, n \in \omega \rangle \\ N^1 &= \langle q_{\alpha n} s_{\alpha n+1}^1 - s_{\alpha n}^1 - \sum_{j < n_\alpha} s_{\alpha_j n_j}^1 - x_{\alpha n} - r_{\alpha n} f : \alpha \in \mathbf{S}, n \in \omega \rangle \end{aligned}$$

Let $F^i = F \oplus C^i/N^i$ and identify $x \in F \oplus C^i$ with $x + N^i$ in F^i . The new relations are

$$(r^i) \quad q_{\alpha n} s_{\alpha n+1}^i = s_{\alpha n}^i + \sum_{j < n_\alpha} s_{\alpha_j n_j}^i + x_{\alpha n} - r_{\alpha n} \delta_{1i} f \quad (i = 0, 1)$$

The maps $\psi_i : F^i \rightarrow H/hR$ given by $\psi_i|_F = 0$, $s_\alpha^i \psi_i = t_\alpha + hR$ and $s_{\alpha n}^i \psi_i = t_{\alpha n} + hR$ preserve the relation in $(*)$ modulo hR and are epic, hence (2) follows.

The proof of (1) is the same as in (3.1).

Suppose $\varphi \in \text{End } F$ extends to $\varphi^i \in \text{End } F^i$ ($i = 0, 1$). Again we consider the pushout $F^0 + F^1 = D$ with $F = F^0 \cap F^1$ as in (3.1), and let

$$U = \langle F, d_{\alpha n}, d_{\alpha'} : \alpha \in \mathbf{S}, n \in \omega, \alpha' \in \mu \setminus \mathbf{S} \rangle \subseteq D$$

for $d_{\alpha n} = s_{\alpha n}^0 - s_{\alpha n}^1$ and $d_\alpha = s_\alpha^0 - s'_\alpha$. We also have $U + F^i = D$ and $U \cap F^i = F$, however the relations for $d_{\alpha n}$ from (r^i) are

$$q_{\alpha n} d_{\alpha n+1} = d_{\alpha n} + \sum_{j < n_\alpha} d_{\alpha_j n_j} + r_{\alpha n} f. \text{ Hence}$$

$$V = \langle f, d_{\alpha n}, d_{\alpha'} : \alpha \in \mathbf{S}, n \in \omega \text{ and } \alpha' \in \mu \setminus \mathbf{S} \rangle \cong H$$

by (*). We find a projection $\pi : D \rightarrow V$, because - as in (3.1) - fR has a complement L_{-1} in $K_0 \subseteq F$ (cf. 3.1) and we find complements L_n of each K_n in K_{n+1} , hence $D \subseteq V \oplus \bigoplus_{n \geq -1} L_n$ gives π . The pushout D provides $\Phi \in \text{End } D$ extending φ^i ($i = 0, 1$) and $\Phi' = \Phi|V \cdot \pi$ is an endomorphism of V . Finally $\text{End } H = R$ forces $f\varphi = f\Phi \in fR$.

Proof (6.1): The proof is similar to §5, however (3.1) is replaced by (6.2). We only indicate the changes. Assuming $\lambda = \mu^+ = 2^\mu$ we prepare the application of $\Phi_\lambda(E)$ for $E = \{\alpha \in \lambda, cf\alpha = \omega\}$. Condition (2) in §5 must be replaced by a new condition

$$(2) \begin{cases} X \text{ is an endomorphism } h_X \in \text{End } G_i. \text{ If } K_j^\omega = \langle x_{\alpha j} : \alpha \in A_{ij} \rangle \\ (j \in a_i) \text{ and if we identify } (G_i, \bigoplus_{j < n} K_j^\omega, g_\beta, h_X : n \in \omega) \text{ with} \\ (F, \bigoplus_{j < n} K'_j, f, \varphi : n \in \omega) \text{ in (6.2), then we} \\ \textbf{require that } \varphi \text{ does not extend to } \text{End } F^0. \end{cases}$$

In the construction of G we must change condition (iii). Now we have

$$(iii) \begin{cases} \text{If } i \in E_\beta \text{ and (1) holds, then we choose } H_i \text{ as an extension} \\ F^{\varphi_\beta(i)} \text{ given by the identification } (F, \bigoplus_{j < n} K'_j, f, \varphi : n \in \omega) \\ \text{in (6.2) with } (G_i, \bigoplus_{j < n} K_j^\omega, g_\beta, h_X : n \in \omega). \end{cases}$$

As in §5, the final R -module G has size λ and is strongly λ -free. In particular $R \subseteq \text{End } G$ by scalar multiplication. If $\psi \in \text{End } G \setminus R$, then we can find a cub

$$C = \{i < \lambda : \psi|G_i = \psi_i \in \text{End } G_i \setminus R\}.$$

Moreover we can choose $f \in G_i$ with G_i/fR free such that $f\psi \notin fR$ and $f = g_\beta$ by enumeration. Choose an ordinal $i \in C \cap \mathcal{X}_\beta(\psi)$ and observe that ψ_i extends to an endomorphism $\psi' \in \text{End } G_{i+1}$. The latter follows from the fact that $G_{i+1}\psi \subseteq G_j$ for some $j < \lambda$ and G_{i+1} is a summand of G_j , so $\psi' = \psi|G_{i+1} \cdot \pi$ for some projection π . Hence $P_i^\beta(\psi_i) = 1$ and $G_{i+1} = F^1$ as in the Step-Lemma 6.2. However $P_i^\beta(\psi_i) = 1$ and (2) tell us that ψ_i extends to F^0 as in (6.2) as well. Therefore $f\psi \in fR$ by (6.1), a contradiction, and $\text{End } G = R$ follows.

Inductively we derive a

Corollary 6.3: *If $2^{\aleph_i} = \aleph_{i+1}$ for $i = 0, \dots, n-1$, then there exists a strongly $\aleph_n$ -free abelian group G with $|G| = \aleph_n$ and $\text{End } G = R$.*

We also have a corollary from the proof of (6.1) observing that the R -algebra R can be replaced by an R -algebra A .

Theorem 6.1*: *Let R be a countable, commutative ring which is S -reduced and S -torsion-free for some multiplicatively closed set S of regular elements in R . Assume $\lambda = 2^\mu = \mu^+$ for some regular cardinal μ and suppose there exists a strongly μ -free A -module H of cardinality μ with $\text{End}_R H = A$ and $|A| < \mu$, A_R free. Then we can find a strongly λ -free A -module G of cardinality λ with $\text{End}_R G = A$.*

Hence we have a corollary which ensures counterexamples for Kaplansky's test problems, cf. Corner [3] and Corner, Göbel [4]. The next corollary strengthens (6.3).

Corollary 6.3*: *Let R be as in (6.1*) and assume $2^{\aleph_i} = \aleph_{i+1}$ for $i = 0, \dots, n-1$. If A is a countable, free R -algebra, then there exists a strongly $\aleph_n$ -free R -module G with $|G| = \aleph_n$ and $\text{End } G = A$.*

§7 Almost free abelian groups which are of cardinality $> \aleph_\omega$

In order to derive results like the existence of strongly $|G|$ -free groups G with $|G| = \aleph_{\omega+1}$ and $\text{End } G = \mathbb{Z}$ under G.C.H., the induction used in §§3, 4 and 5 breaks down at $\aleph_\omega$ and we must extend the basic results of the earlier sections in order to overcome this difficulty. The changes will be sketched below. If μ is a regular cardinal, then let $\mathbf{r}(\mu)$ be the set of all decreasing sequences $u = \langle \mu_1, \dots, \mu_n \rangle$ of regular cardinals $\mu \geq \mu_1 > \dots > \mu_n$. Recall the notion of a u -resolution from (2.3) and of the freeness type of an R -module H from §2. In case $\text{End } H = A$ we assume that H is a strongly μ -free A -module and replace R by A in the above definitions. If H_0/hA is A -free for H_0 as in (2.2) and $h \in H_0$, then $\text{ft } H = \text{ft}(H/hA)$ follows. Observe that we showed in (3.1) (1) and (2) that a strongly μ -free R -module H of freeness type $u = \langle \mu, \aleph_0 \rangle$ has a $(\aleph_0)$ -resolution. This implication does not depend on additional properties of H as $H^* = 0$ or $\text{End } H = A$.

Moreover, we must refine the use of ω -sequences converging to particular ordinals. If $u = \langle \mu_1, \dots, \mu_n \rangle \in \mathbf{r}(\mu)$, then let $\times u = \mu_1 \times \dots \times \mu_n$ be equipped with the lexicographical order. We consider strictly order preserving monomorphisms $\alpha : \times u \rightarrow \mu$ and let $a = \text{Im}(\alpha)$ and $i = \sup a$ which is cofinal to μ_1 . We say that $X \subseteq a$ is bounded (with respect to α) if there is $\beta = (\beta_1, \dots, \beta_n) \in \times u$ such that $X \subseteq \text{Im}(\alpha|_\beta)$.

We may view α as an “ n -ladder” at a . A bounded set X is bounded “on each ladder” in an obvious sense. If $n = 1$, then a is a μ_1 -sequence representing $\text{cf}(i) = \mu_1$. The bounded subsets of a constitute an ideal of the set of subsets

$\mathcal{P}(a)$ of a . The set $a \subseteq i$ will be used to find an extension on $G_i \subseteq G_{i+1}$ which will lead to the desired module $G = \bigcup_{i < \lambda} G_i$. If $u = \langle \aleph_0 \rangle$, as is the case considered in §3-6, then the 1-ladder system is an ω -sequence a_i at each ordinal i with $cf(i) = \omega$ which has an additional property (+) (§3) which ensures freeness of G_δ at limit ordinals $\delta < \mu^+$. A similar condition is required for a general u :

$$(+) \left\{ \begin{array}{l} \text{If } cf(i) = \mu_1, \text{ then there exists an } n\text{-ladder system } a_i \subseteq i \\ \text{such that for each } \delta < \mu^+ \text{ there exist bounded subsets } b_i \subseteq a_i \\ \text{(as defined above) such that the sets } (a_i \setminus b_i) \text{ } i \in S_{\mu_1}, \text{ } i < \delta \\ \text{are pairwise disjoint.} \end{array} \right.$$

This is ensured by the following combinatorial result of Shelah, which follows from the main theorem of [35].

Theorem 7.1 (ZFC + G.C.H.): *If $S = \{i \in \aleph_{\omega+1}, cf(i) = \omega_1\}$ then $\diamond_{\aleph_{\omega+1}}(S)$ holds. Moreover, for any $i \in S$ we can find a 2-ladder system $\alpha^i : \omega_1 \times \omega \rightarrow i$ such that the sets $a_i = Im(\alpha^i)$ satisfy (+).*

The μ - $\aleph_0$ -Step-Lemma 3.1 and (6.2) can be refined to

The μ - u -Step-Lemma 7.2: *Let (R, S) be as in (2.1) and A be a countable, free R -algebra. We will distinguish two cases (a) and (b). In case (a) let $H = H'$ be a strongly μ -free R -module of cardinality μ of freeness type $ftH = u$. In case (b) let H be a strongly μ -free A -module of cardinality μ with $End_R H = A$ and $h \in H_0$ (in (2.2)) such that $ftH = ftH' = u$ for $H' = H/hA$. Then we can find two u -resolutions of H' for μ_1 (with respect to R in case (a) and to A in case (b)) of the form F^j/K ($j = 0, 1$) such that for any basic element $f \in K_0$ the following holds*

- (a) *If $H^* = 0$ and $\varphi \in K^*$ extends to both $\varphi^j \in (F^j)^*$ ($j = 0, 1$), then $f\varphi = 0$.*
- (b) *If $\varphi \in End_R(K)$ extends to both $\varphi^j \in End F^j$ ($j = 0, 1$), then $f\varphi \in fA$.*

Proof. We consider case (a) only and must find suitable extensions F^j of K . Recall that $K = \bigoplus_{i < \mu} K_i$ for $\mu = \mu_1$ is a direct sum of free R -modules K_i of rank μ , cf. (2.3). We will define F^j for $j = 0, 1$. We fix an n -ladder $a \subseteq \mu$ and write K as a continuous increasing chain $K = \bigcup_{i < \mu} F_i$ of free R -modules of rank μ such that $F_{i+1} = F_i \oplus K_i$ for all $i \in a$. The module H can also be represented by (2.2) using a in the form $H = \bigcup_{\alpha \in a} H_\alpha$. Using the ordering on a (induced by μ and also by $\times u$), equations (2.2) can be expressed as

$$q_\alpha t_{\alpha+1} = t_\alpha + \sum_{\alpha_j < \alpha} t_{\alpha_j}.$$

The free resolution is now similar to (3.1)(r^j). Defining C^j and N^j according to (3.1) we let F^j ($j = 0, 1$) be generated by K and new elements s_α^j ($\alpha \in a$) subject to the relations

$$q_\alpha s_{\alpha+1}^j = s_\alpha^j + \sum_{\alpha_j < \alpha} s_{\alpha_j}^j + x_\alpha + \delta_{j1} f,$$

where $x_\alpha \in K_\alpha$ are fixed basic elements. Arguments as in (3.1) will give (a) and (b).

In order to ensure that unions at limit ordinals remain free in the construction of §5, the ω -Freeness-Proposition 3.2 was used. At limit ordinals not cofinal to ω , the construction was trivial. This is different here. If $u = \langle \mu_1, \dots, \mu_n \rangle$ is a certain type of freeness, then the new construction will be non-trivial at limit ordinals cofinal to some μ_i . Hence we have to work at those limit ordinals to ensure freeness of the constructed μ^+ filtration. We need the generalized

u -Freeness-Proposition 7.3: *Let μ be a regular cardinal, $u = \langle \mu_1, \dots, \mu_n \rangle \in \mathbf{r}(\mu)$ some freeness type with $\mu_1 < \mu$ and $\delta < \mu^+$ an ordinal with $\{G_i : i \leq \delta\}$ an ascending, continuous chain of submodules with the following properties for all $i < \delta$.*

- (a) G_i is free of rank μ .
- (b) G_{i+1}/G_i is strongly μ -free and let $\mathbf{S} = \{i < \delta : G_{i+1}/G_i \text{ is not free}\}$.
- (c) There exist $G_i \subseteq H_i \subseteq G_{i+1}$ and H_i^0 a free summand of G_i with free quotient such that $H_i = G_i \oplus_{H_i^0} H_i^1$ is a pushout with $H_i^0 \subseteq H_i^1$ free modules of rank μ .
- (d) $G_{i+1} = H_i \oplus \bigoplus_{\tau \in u} K_i^\tau$ with K_i^τ freely generated by $x_{\alpha i}^\tau$ ($\alpha < \mu$).
- (e) If $i \in \mathbf{S}$, then $G_i = H_i$ and $H_i^1 = H_i^0$ (hence $G_{i+1} = G_i \oplus \bigoplus_{\tau \in u} K_i^\tau$).
- (f) If $i \notin \mathbf{S}$, then $\text{cf}(i) = \kappa \in u$. Moreover
 - (i) there exists $a_i \subseteq i$ of order type κ with $a_i \cap \mathbf{S} = \emptyset$ and $\sup a_i = i$.
 - (ii) there exist $A_{ij} \subseteq \mu$ for all $j \in a_i$ such that $A_{ij} \cap A_{ij'} = \emptyset$ for $j \neq j'$ and $A_{ij} \cap A_{i'j'}$ is bounded in μ for $(ij) \neq (i'j')$.
 - (iii) Let $K_j^\kappa = \langle x_{\alpha j}^\kappa : \alpha \in A_{ij} \rangle$, $H_i^0 = \bigoplus_{j \in a_i} K_j^\kappa$. If $k \in a_i$, $\beta < \mu$ and $K_{\beta k}^\kappa = \bigoplus_{j \in a_i}^{j < k} K_j^\kappa \oplus \langle x_{\alpha j}^\kappa : j \in a_i, j \geq k, \alpha < \beta \rangle$, then $H_i^1/K_{\beta k}^\kappa$ is free.

It follows G_δ is free as well.

The proof of (7.3) follows the proof of (3.2) up to minor changes.

Combining (7.1) and our next result (7.5) we can pass $\aleph_\omega$ and obtain the existence of certain modules of cardinal $\lambda = \aleph_{\omega+1}$.

Corollary 7.4 (ZFC + G.C.H.): *If (R, S) is as in §2 and A is a countable, free R -algebra, then we can find strongly $\aleph_{\omega+1}$ -free R -modules G of cardinality $\aleph_{\omega+1}$ with $\text{End}_R G = A$ and of freeness type $\langle \aleph_{\omega+1}, \aleph_1 \rangle$.*

We would like to remark that a further extension of (7.5) would provide modules as in (7.4) for cardinals $\aleph_{\omega k+1}$ ($k \in \omega$). Clearly (7.4) in conjunction with (5.1) and (6.1*), respectively, provides examples for all $\aleph_{\omega+n}$ ($n \geq 1$). A similar result then follows for all $\aleph_{\omega k+n}$ ($k, n \geq 1$). Magidor and Shelah [31] show, however, that there are models of ZFC + G.C.H. in which $\aleph_{\omega^2+1}$ -free abelian groups of cardinality $\aleph_{\omega^2+1}$ are free. This shows that the results above cannot be extended any further.

Theorem 7.5 (ZFC + G.C.H.): *Suppose that the following conditions hold.*

- (1) *Let (R, S) be as in §2 and A a countable, free R -algebra.*
- (2) *There exists a strongly μ -free A -module H of cardinality μ with $\text{End}_R H = A$ of freeness type $u = \langle \mu_1, \dots, \mu_n \rangle$.*
- (3) *Let λ be a regular, uncountable cardinal with a stationary, non-small subset S of ordinals cofinal to some cardinal μ_1 .*
- (4) *There exists an n -ladder system $a_i \subseteq i$ ($i \in S$) with $(+)$ and $\sup a_i = i$, $a_i \cap S = \emptyset$.*

If $\lambda = \mu^+$, then there exists a strongly λ -free R -module G with $|G| = \lambda$, $\text{ft}(G) = \lambda^\wedge u$ and $\text{End}_R G = A$.

Remark: A similar result holds for modules with trivial dual.

Proof. We indicate the changes in the construction of G , which are based on (2.2) using $u^n = \times u$ and the n -ladder $\alpha^i : \times u \rightarrow i$ with $\text{Im}(\alpha^i) = a_i$. Recall that $\text{cf}(i) = \mu$. From $a_i \cap S = \emptyset$ and the inductive construction of $F = \bigcup_{j < \lambda} F_j$ we have $F_{\alpha+1} = F_\alpha \oplus K_\alpha$ for $\alpha \in a_i$. We choose basic elements $x_\alpha \in K_\alpha$ (with respect to A in case of $\text{End}_R G = A$ and with respect to R for $G^* = 0$) which are used in (7.2) in order to find F_{i+1}^j ($j = 0, 1$) with the desired properties. Moreover, (7.3) takes care of freeness at limit ordinals. The prediction principle (7.5)(c) picks the right branch in 2^λ which leads to $\text{End}_R G = A$ and $G^* = 0$, respectively.

References

- [1] R. BAER: *Abelian groups without elements of finite order*, Duke Math. J. **3** (1937), 68–122.
- [2] A.L.S. CORNER: *Every countable reduced torsion-free ring is an endomorphism ring*, Proc. London Math. Soc. (3) **13** (1963), 687–710.
- [3] A.L.S. CORNER: *Additive categories and a theorem of W.G. Leavitt*, Bull. Amer. Math. Soc. **75** (1969), 78–82.
- [4] A.L.S. CORNER and R. GÖBEL: *Prescribing endomorphism algebras, a unified treatment*, Proc. London Math. Soc. (3) **50**(1985), 447–479.
- [5] A.L.S. CORNER and R. GÖBEL: *On the existence of an $\aleph_1$ -free abelian group of cardinality $\aleph_1$ with no free summands* (unpublished).
- [6] K. DEVLIN and S. SHELAH: *A weak version of $\diamond$ which follows from $2^{\aleph_0} < 2^{\aleph_1}$* , Israel J. Math. **29** (1978), 239–247.
- [7] M. DUGAS: *Fast Freie Abelsche Gruppen mit Endomorphismenring $\mathbb{Z}$* , Journal of Algebra **71** (1981), 314–321.
- [8] M. DUGAS and R. GÖBEL: *Every cotorsion-free ring is an endomorphism ring*, Proc. London Math. Soc (3) **45** (1982), 319–336.
- [9] M. DUGAS and R. GÖBEL: *Every cotorsion-free algebra is an endomorphism algebra*, Math. Zeitschr. **181** (1982), 451–470.
- [10] M. DUGAS and R. GÖBEL: *On radicals and products*, Pacific J. Math. **18** (1985), 70–104.
- [11] K. EDA: *Cardinal restrictions for preradicals*, pp.277–283 in “Abelian Group Theory”, Contemporary Math. **87**, Providence (1989).
- [12] P.C. EKLOF: *On the existence of κ -free abelian groups*, Proc. Amer. Math. Soc. **47** (1975), 65–72.
- [13] P.C. EKLOF: *Set theoretic methods in homological algebra and abelian groups*, Les Presses de l’Université de Montréal, Montreal 1980.
- [14] P.C. EKLOF and A.H. MEKLER: *On constructing indecomposable groups in L* , Journal of Algebra **49** (1977), 96–103.
- [15] P.C. EKLOF and A.H. MEKLER: *Almost free modules, set-theoretic methods*, North-Holland, Amsterdam 1990.
- [16] P.C. EKLOF and S. SHELAH: *On Whitehead modules*, Journal of Algebra **142** (1991), 492–510.

- [17] L. FUCHS: *Infinite abelian groups, Vol. I*, Academic Press, New York, 1970.
- [18] L. FUCHS: *Infinite abelian groups, Vol. II*, Academic Press, New York, 1973.
- [19] R. GÖBEL: *An easy topological construction for realizing endomorphism rings*, Proc. R. Irish Acad. **92 A** (1992) 281-284.
- [20] R. GÖBEL and W. MAY: *Independence in completions and endomorphism algebras*, Forum Mathematicum **1** (1989), 215-226.
- [21] R. GÖBEL and W. MAY: *Four submodules suffice for realizing algebras over commutative rings*, J. Pure Appl. Algebra **65** (1990), 29-43.
- [22] R. GÖBEL and S. SHELAH: *On the existence of rigid $\aleph_1$ -free abelian groups of cardinality $\aleph_1$* , to appear in Proceedings of the Padova Conference on Abelian Groups and Modules 1995
- [23] R. GÖBEL and B. GOLDSMITH: *The Kaplansky Test problem — an approach via radicals*, Journ. Pure and Appl. Algebra **99** (1995), 331-344.
- [24] J. GREGORY: *Higher Souslin trees and the generalized continuum hypothesis*, Journ. Symb. Logic **41** (1976), 663-667.
- [25] PHILLIP GRIFFITH: *$\aleph_n$ -free abelian groups*, Quart. J. Math. (2) **23** (1972), 417-425.
- [26] G. HIGMAN: *Almost free groups*, Proc. London Math. Soc. **1** (1951), 184-190.
- [27] G. HIGMAN: *Some countably free groups*, pp. 129-150 in Proceedings “Group Theory”, Singapore 1991.
- [28] P. HILL: *New criteria for freeness in abelian groups II*, Trans. Amer. Math. Soc. **196** (1974), 191-201.
- [29] T. JECH: *Set Theory*, Academic Press, New York 1978.
- [30] R. JENSEN: *The fine structure of the constructible hierarchy*, Ann. of Math. Logic **4** (1972), 229-308.
- [31] M. MAGIDOR and S. SHELAH: *When does almost free imply free? (for groups, transversals, etc.)*, Journ. Amer. Math. Soc. **7** (4) (1994), 769-830.
- [32] A.H. MEKLER: *How to construct almost free groups*, Can. J. Math. **32** (1980), 1206-1228.
- [33] J. ROTMAN: *Homological Algebra*, Academic Press, New York 1979.

- [34] S. SHELAH *A compactness theorem for singular cardinals, free algebras, Whitehead problem and transversals*, Israel Journal of Math., **21** (1975), 319-349.
- [35] S. SHELAH: *On successors of singular cardinals*, Logic Colloquium '78 **97** (1978) of Stud. Logic Foundations Math., 357-380, North Holland, Amsterdam-New York.
- [36] S. SHELAH: *On uncountable abelian groups*, Israel J. Math. **32** (1979), 311-330.
- [37] S. SHELAH: *Whitehead groups may not be free, even assuming H . II* Israel J. Math. **35** (1980), 257 - 285
- [38] S. SHELAH: *On endo-rigid strongly $\aleph_1$ -free abelian groups in $\aleph_1$* , Israel J. Math., **40** (1981), 291-295.
- [39] S. SHELAH: *A combinatorial principle and endomorphism rings. I: On p -groups*, Israel J. Math. **49** (1984), 239-257.
- [40] S. SHELAH: *A combinatorial theorem and endomorphism rings of abelian groups. II* pp. 37 - 86 in *Abelian groups and modules* CISM Lecture Notes, Vol. **287**, Springer Wien-New York 1984.
- [41] S. SHELAH: *Incompactness in regular cardinals*, Notre Dame Journal of Formal Logic, **26** (1985), 195-228.
- [42] S. SHELAH: *A note on κ -freeness of abelian groups*, pp. 260 - 268 in *Around classification theory of models*, Lecture Notes in Math. Vol. **1182**, Springer, Berlin, 1986.
- [43] S. SHELAH and Z. SPASOJEVIC: *A forcing axiom on strongly inaccessible cardinals making all uniformizations for all ladder systems lying on a fixed stationary set and applications for abelian groups*, manuscript No. 587.